



Amphibians and reptiles of the Ankaratra Massif: reproductive diversity, biogeography and conservation of a montane fauna in Madagascar

MIGUEL VENCES

Institute for Biodiversity and Ecosystem Dynamics, Zoological Museum, University of Amsterdam, P.O. Box 94766, 1090 GT Amsterdam (The Netherlands)
E-mail: m.vences@online.de

FRANCO ANDREONE

Museo Regionale di Scienze Naturali, Via G. Giolitti 36, I-10123 Torino (Italy)
E-mail: f.andreone@libero.it

FRANK GLAW

Zoologische Staatssammlung, Münchenstr. 21, D-81247 München (Germany)
E-mail: Frank.Glaw@zsm.mun.de

NOROMALALA RAMINOSOA

Université d'Antananarivo, Département de Biologie Animale, Antananarivo (101) (Madagascar)
E-mail: rsnor@yahoo.fr

JASMIN EMILE RANDRIANIRINA

Parc Botanique et Zoologique de Tsimbazaza, Département Faune BP 4096, and Ecology Training Programme BP 738, Antananarivo (101) (Madagascar)
E-mail: jbza@mts.mg

DAVID RODRIGUEZ VIEITES

Departamento de Biología Animal, Laboratorio de Anatomía Animal, Buzón 137, Facultad de Ciencias Biológicas, Universidade de Vigo, Apdo. 874, E-36200 Vigo (PO) Galicia (Spain)
E-mail: dvieites@univigo.es

ABSTRACT

The Ankaratra Massif in central Madagascar includes rainforest between ca. 1500 and 2000 m elevation and montane savannah and heathland between ca. 2000 and 2642 m elevation. It is populated by 15 species of amphibians and 12 species of reptiles, all of which were observed in our surveys between 1992-2001. Additionally, the Ankaratra material housed in several herpetological collections was revised. Examination of type material showed that *Scaphiophryne madagascariensis* is the name to be applied to the scaphiophryine microhylid frog occurring at Ankaratra; the previously used name *S. pustulosa* is a junior synonym, while populations from the Andringitra Massif so far considered as *S. madagascariensis* do not belong to this species. Reproductive behaviour and advertisement calls are described for *S. madagascariensis*. Two other frog species (*Boophis williamsi*, *Mantidactylus pauliani*) were observed for the first time for almost 30 years; their live coloration and morphology are described. The skink *Amphiglossus macrocerus* is shown to be live-bearing. Breeding in the austral summer could be demonstrated for most species, including the geckos *Lygodactylus mirabilis* and *Phebuma barbouri*

(Received 16 May 2001 - Accepted 22 February 2002)

which are known to lay eggs also in the cold winter months (July-September). The higher-level taxonomic composition of the Ankaratra herpetofauna recalls that of the rainforests of eastern Madagascar, and also the species show mainly eastern affinities. A number of groups (especially snakes) are under-represented and some (e.g., nocturnal geckos and dwarf chameleons) are totally lacking. Three high-elevation specialists (*B. williamsi*, *M. pauliani*, *L. mirabilis*) appear to be Ankaratra endemics (11% of the overall community). Pending further surveys and taxonomic revisions, more than one fifth of the herpetofauna (six species or more) may be endemic or semi-endemic to the region, which is not included in the network of Madagascar's protected areas. A monitoring program is therefore proposed to assess the conservation status of these poorly known taxa.

KEY WORDS: Amphibia - Reptilia - Madagascar - Biodiversity - Ankaratra Massif - Montane heathland - Rainforest - Conservation.

ACKNOWLEDGEMENTS

We are grateful to G. Besmer (Ambatolampy) and J. Gräbener (Antananarivo) of the Gesellschaft für Technische Zusammenarbeit (GTZ), as well as to the teams of the GTZ projects PDFIV and POLFOR and the local staff of the Ministère des Eaux et Forêts, for the logistic and technical support granted during the surveys. In the period 1-5.III.2000, we were aided in the field by V. A. Rasoaampiarina, E. Doumenq, H. V. Razakarivony, A. R. Andrianajivoarivelo, C. Ramamontisoa, H. E. L. Rafalimanantsoa and D. Andriafidison, and in 2001 by D. Rakoromalala and L. Raharivoloniaina. In previous surveys F. Andreone and J. E. Randrianirina were accompanied by D. Vallan and S. De Michelis. Other specimens now housed in MRSN were collected by R. Nincheri. The chameleon survey at Analamilona was carried out as "Projet Expérimental" of the Ministère des Eaux et Forêts (Antananarivo-Madagascar), in collaboration with the "Langaha" Association. A. Dubois and A. Ohler (MNHN, Paris), W. Böhme (ZFMK, Bonn), B. T. Clarke, E. N. Arnold and C. McCarthy (BMNH, London), and B. von Tuijl (ZMA, Amsterdam) allowed examination of specimens held in their care. R. Blommers-Schlösser made call recordings available. We also wish to thank L. Luiselli and an anonymous referee for their helpful comments upon a former version of this manuscript, as well as N. E. Baldacini for the editorial assistance.

INTRODUCTION

Madagascar, the fourth largest island in the world, has a varied topography, with a central chain spanning from north to south and including peaks as high as 2876 m (Tsaratanana), 2133 m (Marojejy), 2642 m (Tsi-afajavona, Ankaratra Massif), 2658 m (Pic Boby, Andringitra Massif), and 1956 m (Anosy Massif) a.s.l. These high-elevation regions of Madagascar are populated by amphibian and reptile communities partly specialized to montane habitats. Several of these species are endemic to the massifs or to montane areas (Raxworthy & Nussbaum, 1996a; Vences & Glaw, 1999). While Andringitra, Anosy and Marojejy have been subject of extensive herpetofaunistic surveys in the past decade (Raxworthy & Nussbaum, 1996a, b; Andreone & Randriamantazo, 1997; Raxworthy *et al.*, 1998; Nussbaum *et al.*, 1999; Raselimanana, 1999; Raselimanana *et al.*, 2000), no recent comprehensive work is available on the amphibians

and reptiles of the easily accessible Ankaratra mountains south of Madagascar's capital Antananarivo. This may be due to two reasons. First, only small remains of primary (partly degraded) forest are found at Ankaratra, almost completely concentrated in the area managed by the Manjakatempo forestry station; second, due to the easy access to Manjakatempo and Ankaratra in general, many isolated works have been conducted on the herpetofauna of the region (Guibé, 1952; Pasteur, 1959, 1962; Andreone, 1993; Glaw & Vences, 1994), which leaves few 'spectacular' discoveries to be expected during intensive inventories.

The present paper reports the results of several surveys conducted during 1992, 1994, 2000 and 2001 in the Manjakatempo, Nosiarivo, and Tsiarafajavona areas. The distribution and biological information available on the amphibians and reptiles of Ankaratra is summarized, and their biogeographic relationships and conservation status are discussed.

MATERIALS AND METHODS

Methods of analysis of preserved materials

The present data are based on the examination of specimens housed in the Natural History Museum (formerly British Museum of Natural History; BMNH), Muséum National d'Histoire Naturelle, Paris (MNHN), Museo Regionale di Scienze Naturali, Torino (MRSN and MRSN-FAZC), Université d'Antananarivo, Département de Biologie Animale (UADBA and RD), Zoologisches Forschungsinstitut und Museum A. Koenig, Bonn (ZFMK), Zoologisch Museum Amsterdam (ZMA), and Zoologische Staatssammlung, München (ZSM). UADBA specimens are given with preliminary numbers which correspond to the provisional field numbers assigned by F. Glaw and M. Vences (FG/MV) in 2000 and the ones assigned by M. Vences (MV) in 2001. For each species, a mention is made of the Ankaratra voucher specimens housed in these collections. When necessary for morphological, taxonomical or biogeographical discussions, also the specimens examined from other localities are listed as "further material".

Morphometric measurements were carried out by the senior Author with a calliper to the nearest tenth of a millimetre. SVL is used as abbreviation for snout-vent length. Bioacoustic analyses were made by F. G. with the sound analysing system MEDAV-Spektro 3.2. Bioacoustic and morphometric values are generally presented as range (mean ± SD in parentheses).

Acronyms for the quoted collectors and authors are as follows: CPB, C. P. Blanc; DR, D. Rakotomalala; DRV, D. R. Vieies; DV, D. Vallan; FA, F. Andreone; FG, F. Glaw; JER, J. E. Randrianirina; MV, M. Vences; NR, N. Raminosoa; RN, R. Nincheri; SDM, S. De Michelis.

Study area

Geologically (Perrier de la Bâtie, 1927; Besairie, 1954), the Ankaratra is an important and slightly curved barrier roughly extending in a north-south orientation (Fig. 1). Part of the central watershed that separates the rivers flowing towards the Mozambique channel from those flowing into the Indian Ocean, it is made up by different, partly overlapping volcanic zones which laterally extend about 30 km, contacting neighbouring volcanic centres. The northern part of the massif mostly corresponds to a younger, Pleistocene volcanic activity, with characteristic blackish, compact lavas. This area includes the highest peaks: Tsiarafajavona (2642 m), Tsiarafakao (2530 m), Ankavitra (2603 m), Ambohimainty (2595 m), and Ambohimirandrana (2412 m). The southern part, in contrast, originating from older (Pliocene) volcanic activity, is partly characterized by trachytic domes such as that of Famoizankova (2362 m).

The climate in this massif follows the general seasonal pattern typical for Madagascar, with a cold and dry season in the austral winter and a warm and wet one in the austral summer. According to meteorological data recorded at Manjakatempo (Donque, 1975), the mean temperatures range from 11.9° C (July) to 17.9° C (January); the extremes of the mean minimum and maximum temperatures per month were 5.5° C (July) and 24.7° C (November). However, at the high mountain areas of the Tsiarafajavona, low temperatures are also recorded in the austral summer (we recorded less than 10° C at night in February), and temperatures below 0° C are common in the austral winter. Mean annual precipitation is 2012 mm, with a recorded range of 1445-2662 mm. The highest amount of rainfall is in January (362 mm), while the minimum corresponds to June or July, with a mean total of about 200 mm in the whole period May-September.

The vegetation of the massif contains remnants of primary rainforest which are mainly located along the eastern slopes at elevations of 1700-2000 m (Fig. 2a). As discussed by Goodman *et al.*




Fig. 1 - Map (a) of Madagascar, showing areas between 1500-2000 m and above 2000 m elevation and (b) of the Ankaratra region between Ambatolampy and the Tsiarafajavona, showing areas of 2000-2500 m and above 2500 m, forested areas, and main roads. Numbers refer to several of the collection sites listed in Materials and Methods: 1, Manjakatempo; 3, Manjavona Valley; 4, Ambohimirandrana; 5, Nosiarivo; 8, Tsiarafajavona road; 9, Tsiarafajavona Plateau; 10, Tsiarafajavona. The Ankaratra map is a simplified scheme based on the maps of the Malagasy geographical institute at a scale of 1:500,000; positioning of localities is only approximative. For detailed coordinates, see Materials and Methods.

(1996), the forest in the area of the sacred royal Tankarava tombs (Nosiarivo) is almost exclusively composed of a single tree species (*Weinmannia bojeriana*) and may be a historical plantation dating back to former centuries. The rainforest remnants (including the Nosiarivo Forest) are largely concentrated in the area of the Manjakatempo Forest station, which also contains large pine plantations. Roughly above 2000 m, the forests are replaced by an extensive grassland area, with plots of heathland (Fig. 2b) and remains of other vegetation types on steep slopes along brooks. This savannah landscape, with only rare occurrence of higher ericoid bushes, was already present in 1929 (Rand, 1936).

Ankaratra localities reported in the following accounts are (see Fig. 1): 1, Manjakatempo (19°21' S, 47°18' E, ca. 1700 m); 2, Lac

Froid; 3, Manjavona Valley (here considered as corresponding to "vallée Mahavona" in the MNHN catalogue and "Mihavona" in Blommers-Schlösser & Blanc (1991): 19°21' S, 47°16' E, ca. 1800 m); 4, Ambohimirandrana (19°20' S, 47°16' E, ca. 2000 m); 5, Nosiarivo (19°20' S, 47°17' E, ca. 2000 m); 6, Marizana swamp (listed as "marais Marizana" in the MNHN catalogue); 7, Hctay Forest; 8, Tsiarafajavona road (used for different sites along a section of road that leads from the Ambohimirandrana area towards the Tsiarafajavona summit); 9, Tsiarafajavona Plateau (19°20' S, 47°14' E, 2380 m); 10, Tsiarafajavona; 11, Analamilona (19°20' S, 47°18' E, 1800 m). Although not strictly part of the Ankaratra massif, records from 12, Ambatolampy (19°22' S, 47°26' E, 1600 m), are also considered. As Ankaratra slopes are often rather steep, and




Fig. 2 - Ankaratra landscapes, and rediscovered endemic montane frogs: a, Nosiarivo Forest; b, savannah with ericoid vegetation alongside the Tsiarafajavona road; c, d, dorsolateral and ventral view of *Boophis williamsi* from Ambohimirandrana (male, ZSM 734/2001); e, f, dorsolateral and ventral views of *Mantidactylus pauliani* from Ambohimirandrana (ZSM 756/2001).

